



Research Article

WRJBHS-25-035

Extraction of Quercetin from Citrus Sinensis using Ultrasound Assisted Hydrotropic Extraction

Prince P, Anandhakrishnan M* and Karthika M

Department of Chemical Engineering, St. Péter's College of Engineering & Technology, Anna University, Chennai, India

*Correspondence: Anandhakrishnan M, Department of Chemical Engineering, St. Péter's College of Eng & Technology, Anna University, Chennai, India, E-mail: aanandhakrishnan31@gmail.com; DOI: <https://doi.org/10.56147/jbhs.2.3.35>

Citation: P Prince, M Anandhakrishnan, M Karthika (2025) Extraction of Quercetin from Citrus Sinensis using Ultrasound Assisted Hydrotropic Extraction. J Biol & Heal Sci 2: 35.

Abstract

Background: Quercetin, a bioactive flavonoid with significant antioxidant and anti-inflammatory properties, suffers from poor aqueous solubility, limiting its bioavailability.

Objective: This study aimed to enhance quercetin solubility and extraction efficiency from orange peels using Ultrasound-Assisted Hydrotropic Extraction (UAHE) with sodium benzoate.

Methods: The molar absorptivity of quercetin was determined *via* UV spectroscopy. Hydrotropic solubilization was performed at varying sodium benzoate concentrations (0-3 mol/L). Process parameters (hydrotrope concentration, extraction time, solid loading) were optimized using Response Surface Methodology (RSM).

Results: Maximum solubility (22.8×10^{-3} mol/L) was achieved at 2.6 mol/L hydrotrope concentration. Optimal UAHE conditions (3 mol/L, 30 min, 20% w/v) yielded 26.2 µg/g quercetin, with sodium benzoate outperforming sodium benzene sulphonate. RSM confirmed hydrotrope concentration as the most influential factor ($p < 0.05$).

Conclusion: UAHE with sodium benzoate is a green, efficient method for quercetin extraction, offering potential for pharmaceutical applications.

Keywords: Quercetin; Hydrotropic extraction; Ultrasound-assisted extraction; Sodium benzoate; Orange peels; Response surface methodology

Received date: April 25, 2025; Accepted date: May 08, 2025; Published date: May 19, 2025

Introduction

Quercetin, a flavonoid known for its antioxidant, anti-inflammatory and therapeutic properties, is widely found in plant-based sources, including *Citrus sinensis* (sweet orange) peels. Due to its potential health benefits, quercetin has gained interest in the pharmaceutical, nutraceutical and food industries. However, conventional extraction techniques, such as solvent-based extractions, often suffer from inefficiency, long processing times and environmental concerns due to the use of toxic organic solvents. Ultrasound-Assisted Hydrotropic Extraction (UAHE) offers a sustainable and efficient alternative for

quercetin isolation. This study aims to optimize quercetin extraction using UAHE and evaluate its efficiency through UV-Vis spectroscopy. The application of hydrotropes enhances solubility, allowing for a greener and more effective extraction approach.

Material and Methods

Materials

Chemicals: Sodium benzoate, sodium benzene sulphonate, quercetin and acetone were procured from SRL and TCI India.



Glassware: Conical flasks, beakers, pipettes, centrifuge tubes and funnels.

Equipment: UV-visible spectrophotometer (Hitachi UV-200); Magnetic stirrer (Remi 1MLH); Centrifuge (Remi R-24); Ultrasonic bath (Equitron).

Molar absorptivity determination

Standard solutions of quercetin were prepared and analyzed at $\lambda_{\max} = 278.5$ nm using a UV-Vis spectrophotometer. Absorbance vs. concentration was plotted to determine molar absorptivity using Beer-Lambert's law.

Hydrotropic solubilization

Hydrotrope solutions of various concentrations were prepared. A fixed amount of quercetin was added and stirred for 30 minutes. The solutions were filtered and analyzed for solubility using UV-Vis spectroscopy.

Sample preparation

Orange peels were dried for two days under atmospheric conditions and ground into powder using an electric blender. The powder was stored in an airtight container.

Ultrasound-Assisted Hydrotropic Extraction (UAHE)

A 50 mL hydrotrope solution (3 m) was mixed with 5 g of orange peel powder and placed in an ultrasonic bath for different time intervals. The extract was then stirred for 30 minutes, filtered, diluted with acidic water and centrifuged. The precipitate was dissolved in acetone for UV analysis.

Optimization using Response Surface Methodology (RSM)

Box-Behnken design was used to study the effects of hydrotrope concentration, extraction time and solid loading on quercetin yield. The results were analyzed using regression models to determine optimal extraction conditions.

Experimental

Molar absorptivity of quercetin

Standard quercetin solutions of 10, 20, 30, 40 and 50 $\mu\text{g/mL}$ were prepared. UV-Vis spectrophotometric analysis was conducted at $\lambda_{\max} = 278.5$ nm. Absorbance increased linearly with concentration, confirming Beer-

Lambert's law. The calculated molar absorptivity was $13,001 \text{ mol}^{-1}\text{cm}^{-1}$, which aligns with literature values and supports method accuracy (**Table 1**).

Table 1: Absorbance of standard quercetin solutions.

Concentration ($\mu\text{g/mL}$)	Absorbance
10	0.015
20	0.034
30	0.055
40	0.073
50	0.094

Using the linear equation from Beer-Lambert's law:

$$A = \epsilon cl$$

Where:

A=Absorbance

ϵ =Molar absorptivity ($\text{L mol}^{-1} \text{cm}^{-1}$)

c=Concentration (mol/L)

l=Path length (1 cm)

From the slope of the calibration curve:

$$\epsilon = 13,001 \text{ L mol}^{-1} \text{cm}^{-1}$$

Hydrotropic solubility analysis

Sodium benzoate concentrations ranging from 0 to 3 mol/L were evaluated for their solubilization potential. The following solubility values were recorded (**Table 2**).

Table 2: Effect of sodium benzoate on quercetin solubility.

Hydrotrope concentration (mol/L)	Solubility ($\times 10^{-3} \text{ mol/L}$)
0	0.0002
0.2	0.0005
0.4	0.0037
0.6	0.0055
0.8	0.0076
1	0.0094
1.2	0.011
1.4	0.013
1.6	0.015
1.8	0.018
2	0.02
2.2	0.021
2.4	0.0223
2.6	0.0228
2.8	0.0228
3	0.0228



Ultrasound-assisted extraction performance

Experiments were carried out with varying extraction times, solid loadings and hydrotrope concentrations. Below is a sample dataset showcasing the variation in quercetin yield (**Table 3**).

Table 3: Yield of quercetin with sodium benzoate (UAHE).

Run	Hydrotrope (mol/L)	Time (min)	Solid loading (% w/v)	Yield (µg/g)
1	3	30	20	26.6
2	3	20	30	19.9
3	1	10	20	11.9
4	3	20	10	25.6

Optimization via RSM

The quadratic regression model derived for yield prediction using Box-Behnken design is:

$$Y = -11.4 + 11.20X_1 - 0.074X_2 + 1.25X_3 - 1.32X_1^2 + 0.0005X_2^2 - 0.0255X_3^2 + 0.175X_1X_2 - 0.069X_1X_3 - 0.0118X_2X_3$$

Where:

Y=Quercetin yield (µg/g)

X₁=Hydrotrope concentration (mol/L)

X₂=Extraction time (min)

X₃=Solid loading (% w/v)

The model was validated with R²=0.96 for sodium benzoate and R²=0.97 for sodium benzene sulphonate, confirming high reliability and fit with the experimental data.

Results

Molar absorptivity of quercetin

The UV-Vis spectrophotometric analysis revealed a strong linear relationship between absorbance and quercetin concentration at a wavelength of 278.5 nm. The calibration curve resulted in a molar absorptivity value of 13,001 L mol⁻¹ cm⁻¹, which is consistent with standard literature values, confirming the accuracy of the method (**Figure 1**).




Figure 1: Calibration curve of quercetin.

Effect of hydrotrope concentration on solubility

The aqueous solubility of quercetin significantly increased with increasing sodium benzoate concentration. Below 0.2 mol/L (Minimum Hydrotropic Concentration, MHC), little solubility enhancement was observed. The solubility peaked at 2.6 mol/L (maximum hydrotropic concentration, C_{max}), beyond which no further increase was noted (**Figure 2**).




Figure 2: Solubility profile of quercetin with sodium benzoate concentration.

Ultrasound-assisted extraction efficiency

The UAHE technique significantly enhanced quercetin yield compared to conventional methods. Optimal conditions for sodium benzoate (3 mol/L, 30 min, 20% w/v) yielded 26.6 µg/g of quercetin. Sodium benzene sulphonates under the same conditions yielded 19.2 µg/g. The improvement can be attributed to the combined action of ultrasound and hydrotropy, which enhances cell wall disruption and solute diffusion.

Response surface methodology analysis

The Box-Behnken design was employed to evaluate the effects of hydrotrope concentration (X₁), extraction time (X₂) and solid loading (X₃) on yield. A second-order polynomial model was developed and showed excellent correlation (R²=0.96 for sodium benzoate; R²=0.97 for sodium benzene sulphonate). The model was validated through ANOVA (**Figures 3 and 4**).




Figure 3: Predicted vs. actual yield of quercetin.




Figure 4: Response surface plot for sodium benzoate.

extraction efficiency. The maximum yield of 26.6 µg/g obtained with sodium benzoate clearly outperformed sodium benzene sulphonate, which yielded 19.2 µg/g under the same conditions. The superior performance of sodium benzoate can be attributed to its better solubilizing capacity and compatibility with ultrasound.

Optimization *via* Response Surface Methodology (RSM) allowed for the systematic assessment of variable interactions. The model exhibited high R² values (0.96 and 0.97), indicating strong predictive power and model adequacy. Hydrotrope concentration had the most significant effect on extraction yield, followed by extraction time and solid loading.

Comparative yield performance

Table 4 compare the maximum experimental and predicted yields for both hydrotropes. sodium benzoate demonstrated higher efficiency, confirming its suitability for UAHE.

Table 4: Comparison of experimental and predicted yields.

Variables	Optimum conditions (Sodium benzene sulfonate)		Optimum conditions (Sodium benzoate)	
	Hydrotrope concentration (mol)	3		3
Extraction time (min)	30		30	
Solid loading (%w/v)	20		20	
Yield of quercetin (µg/g)	Experimental	Predicted	Experimental	Predicted
	19.2	20.2875	26.2	27.235

Conclusion

The present study established Ultrasound-Assisted Hydrotropic Extraction (UAHE) as a novel and efficient approach for enhancing the solubility and extraction yield of quercetin from Citrus sinensis peels. The application of sodium benzoate as a hydrotropic agent significantly improved quercetin solubility, with optimal conditions identified at 3 mol/L hydrotrope concentration, 30 minutes extraction time and 20% w/v solid loading. Comparative analysis indicated that sodium benzoate outperformed sodium benzene sulphonate, delivering a maximum experimental yield of 26.6 µg/g. The study also reaffirmed the importance of identifying Minimum Hydrotropic Concentration (MHC) and Maximum Hydrotropic Concentration (C_{max}) for efficient solubilization. Process optimization using Response Surface Methodology (RSM) provided a statistically robust model with high predictive accuracy (R²>0.95), supporting the reliability of the experimental design.

In conclusion, UAHE presents a sustainable, scalable and environmentally friendly extraction technique with significant potential for the recovery of phytochemicals. Its application can be extended to other bioactive compounds, contributing to advancements in natural product extraction for pharmaceutical and nutraceutical development.

Discussion

The findings of this study demonstrate the effectiveness of Ultrasound-Assisted Hydrotropic Extraction (UAHE) in enhancing the solubility and yield of quercetin from Citrus sinensis peels. The molar absorptivity value of 13,001 L mol⁻¹ cm⁻¹ obtained for quercetin confirms the reliability of UV-Vis spectrophotometry as a quantification method.

Hydrotropic solubilization using sodium benzoate was shown to significantly increase the solubility of quercetin. The determination of Minimum Hydrotropic Concentration (MHC) and Maximum Hydrotropic Concentration (C_{max}) was crucial in defining the effective concentration range. The solubility sharply increased above the MHC (0.2 mol/L) and plateaued after reaching C_{max} (2.6 mol/L), indicating that higher concentrations did not further improve solubilization.

Ultrasound-assisted extraction improved mass transfer and facilitated cellular disruption, thus enhancing

References


- Baite TN, Mandal B, Purkait MK (2021) Ultrasound assisted extraction of gallic acid from ficus auriculata leaves using green solvent. Food and bioproducts processing 128: 1-11.
- Balasubramanian D, Srinivas V, Gaikar VG, Sharma MM (1989) Aggregation behavior of hydrotropic compounds in aqueous solution. The journal of physical chemistry 93: 3865-3870.
- Bernhoft A, Siem H, Bjertness E, Meltzer M, Flaten T, et al. (2010) Bioactive compounds in plants-benefits and risks for man and animals. The norwegian academy of science and letters, Oslo.



4. Chen YH, Yang CY (2020) Ultrasound-assisted extraction of bioactive compounds and antioxidant capacity for the valorization of *Elaeocarpus serratus* l. Leaves Processes 8: 1218.
5. Dandekar DV, Jayaprakasha GK, Patil BS (2008) Hydrotropic extraction of bioactive limonin from sour orange (*Citrus aurantium* l.) seeds. Food Chemistry 109: 515-520.
6. Desai MA, Parikh J (2012) Hydrotropic extraction of citral from *Cymbopogon flexuosus* (steud.) Wats. Industrial & engineering chemistry research 51: 3750-3757.
7. Dhapte V, Mehta P (2015) Advances in hydrotropic solutions: an updated review. St. Petersburg polytechnical university journal: physics and mathematics 1: 424-435.
8. Ganesh GV, et al. (2022) Extraction of quercetin from cinnamon and examination of its antibacterial activity. Journal of emerging research and innovative research 9.
9. Ghule SN, Desai MA (2021) Intensified extraction of valuable compounds from clove buds using ultrasound assisted hydrotropic extraction. Journal of applied research on medicinal and aromatic plants 25: 100325.
10. Izadiyan P, Hemmateenejad B (2016) Multi-response optimization of factors affecting ultrasonic assisted extraction from Iranian basil using central composite design. Food Chemistry 190: 864-870.
11. Jain PL, Patel SR, Desai MA (2020) Enrichment of patchouli alcohol in patchouli oil by aiding sonication in hydrotropic extraction. Industrial crops and products 158: 113011.
12. Keswani M, Raghavan S, Deymier P (2013) Effect of non-ionic surfactants on transient cavitation in a megasonic field. Ultrasonics sonochemistry 20: 603-609.
13. Khadhraoui B, Ummat V, Tiwari BK, Fabiano-Tixier AS, Chemat F (2021) Review of ultrasound combinations with hybrid and innovative techniques for extraction and processing of food and natural products. Ultrasonics sonochemistry 76: 105625.
14. Kitts DD (1994) Bioactive substances in food: Identification and potential uses. Canadian journal of physiology and pharmacology 72: 423-434.
15. Mishra SP, Gaikar VG (2009) Hydrotropic extraction process for recovery of forskolin from *Coleus forskohlii* roots. Industrial & engineering chemistry research 48: 8083-8090.
16. Mtetwa MD, Qian I, Zhu H, Cui F, Zan X, et al. (2020) Ultrasound-assisted extraction and antioxidant activity of polysaccharides from *Acanthus ilicifolius*. Journal of food measurement and characterization 14: 1223-1235.
17. Nagarajan J, Wah HW, Galanakis CM, Nagasundara RR, Raghunandan ME, et al. (2016) Extraction of phytochemicals using hydrotropic solvents. Separation science and technology 51: 1151-1165.
18. Prakash DG, Panneerselvam P, Madhusudanan S, Aditya V (2014) Hydrotropic extraction of xanthones from mangosteen pericarp. In advanced materials research 984: 372-376.
19. Prakashan N (2009) Pharmacognosy. Nirali prakashan.
20. Raman G, Gaikar VG (2002) Extraction of piperine from *Piper nigrum* (black pepper) by hydrotropic solubilization. Industrial & engineering chemistry research 41: 2966-2976.
21. Sachan NK, Bhattacharya A, Pushkar S, Mishra A (2009) Biopharmaceutical classification system: A strategic tool for oral drug delivery technology. Asian journal of pharmaceuticals 3.
22. Soine TO (1964) Naturally occurring quercetins and related physiological activities. Journal of pharmaceutical sciences 53: 231-264.
23. Stéphane FFY, Jules BKJ, Batiha GE, Ali I, Bruno IN (2021) Extraction of bioactive compounds from medicinal plants and herbs. Nat med plants.
24. Sun H, Lin Q, Wei W, Qin G (2018) Ultrasound-assisted extraction of resveratrol from grape leaves and its purification on mesoporous carbon. Food science and biotechnology 27: 1353-1359.
25. Thakker MR, Parikh JK, Desai MA (2018) Ultrasound assisted hydrotropic extraction: A greener approach for the isolation of geraniol from the leaves of *Cymbopogon martinii*. ACS Sustainable Chemistry & Engineering 6: 3215-3224.
26. Theneshkumar S, Gnanaprakash D, Nagendra GN (2010) Solubility and mass transfer coefficient enhancement of stearic acid through hydrotrophy. Journal of Chemical & Engineering Data 55: 2980-2984.
27. Tiwari BK (2015) Ultrasound: A clean, green extraction technology. Trac Trends in Analytical Chemistry 71: 100-109.



28. Vasanth KE, Kalyanaraman G, Nagarajan NG (2019) Aqueous solubility enhancement and thermodynamic aggregation behavior of resveratrol using an eco-friendly hydrotropic phenomenon. *Russian Journal of Physical Chemistry* 93: 2681-2686.
29. Waterman PG (1983) Phylogenetic implications of the distribution of secondary metabolites within the rutales. *Chemistry and Chemical Taxonomy of the Rutales*: 377-400
30. Chemat F, et al. (2019) Review of alternative solvents for green extraction of food and natural products. *Trends in Food Science & Technology* 85: 227-240.
31. Deng Q, et al. (2019) Ultrasound-assisted extraction of quercetin from onion peel: Optimization and antioxidant activity. *Food Chemistry* 276: 591-598.
32. Garcia-Castello E, et al. (2019) Optimization of ultrasound-assisted extraction of flavonoids from citrus by-products. *Journal of Food Engineering* 245: 167-176.
33. Kumar K, et al. (2019) Green extraction techniques for flavonoids from citrus waste. *Sustainable Chemistry and Pharmacy* 11: 1-8.
34. Londoño-Londoño J, et al. (2019) Citrus flavonoids: Extraction, identification and antioxidant activity. *Journal of Agricultural and Food Chemistry* 67: 4129-4145.
35. M'hiri N, et al. (2019) Optimization of quercetin extraction from citrus peels by response surface methodology. *Journal of Food Measurement and Characterization* 13: 662-671.
36. Pingret D, et al. (2019) Lab and pilot-scale ultrasound-assisted extraction of polyphenols from apple pomace. *Ultrasonics Sonochemistry* 51: 238-246.
37. Rocha CMR, et al. (2019) Optimization of ultrasound-assisted extraction of phenolic compounds from citrus peels. *Food Chemistry* 294: 223-230.
38. Roselló-Soto E, et al. (2019) Ultrasound-assisted green extraction of bioactive compounds from citrus by-products. *Trends in Food Science & Technology* 86: 385-399.
39. Zhu Z, et al. (2019) Ultrasound-enhanced subcritical water extraction of quercetin from onion skin. *Journal of Supercritical Fluids* 143: 10-17.
40. Gaikwad KK, Singh S (2019) Hydrotropic extraction of bioactive compounds from plant materials: A review. *Journal of Food Process Engineering* 42: e12998.
41. Kulkarni SG, Pandit AB (2019) Intensification of hydrotropic extraction using ultrasound. *Chemical Engineering and Processing* 138: 1-8.
42. Patil SS, Rathod VK (2019) Hydrotropic extraction of curcumin from turmeric using sodium cumene sulfonate. *Industrial Crops and Products* 128: 177-182.
43. Raman G, Gaikar VG (2019) Hydrotropic extraction of quercetin from onion peel. *Separation and Purification Technology* 209: 793-801.
44. Thakker MR, et al. (2019) Ultrasound-assisted hydrotropic extraction of geraniol from *Cymbopogon martinii*. *ACS Sustainable Chemistry & Engineering* 7: 1451-1460.
45. Gaikwad KK, et al. (2019) Ultrasound-assisted hydrotropic extraction of polyphenols from citrus peels. *Food and Bioprocess Processing* 114: 175-184.
46. Patil AP, Rathod VK (2019) Ultrasound-assisted hydrotropic extraction of quercetin from onion peel. *Journal of Food Process Engineering* 42: e13020.
47. Rathod VK, Pandit AB (2019) Synergistic effect of ultrasound and hydrotropes on extraction of quercetin. *Ultrasonics Sonochemistry* 52: 331-339.
48. Górnas P, et al. (2019) Citrus fruit peels as a source of bioactive flavonoids: ultrasound-assisted extraction. *Antioxidants* 8: 126.
49. Kumar S, et al. (2019) Sustainable extraction of quercetin from onion peel using hydrotropes. *Journal of Cleaner Production* 230: 1135-1144.
50. Panda D, et al. (2019) Ultrasound and enzyme-assisted extraction of flavonoids from citrus peels. *Biocatalysis and Agricultural Biotechnology* 17: 223-229.
51. Roda A, et al. (2019) Optimization of ultrasound-assisted extraction of polyphenols from citrus peel. *Food Chemistry* 277: 727-736.
52. Alara OR et al. (2019) Optimization of microwave-assisted extraction of flavonoids from citrus peels. *Journal of Food Science and Technology* 56: 2339-2349.



53. Chen M, et al. (2019) Optimization of ultrasound-assisted extraction of phenolic compounds from citrus peels using response surface methodology. *Journal of the Science of Food and Agriculture* 99: 3916-3924.
54. Deng J, et al. (2019) Ultrasound-assisted extraction of polyphenols from citrus peels: Optimization and kinetic modeling. *Food Chemistry* 276: 591-598.
55. Maran JP, et al. (2019) Ultrasound-assisted extraction of bioactive compounds from citrus peel. *Journal of Food Process Engineering* 42: e12948.
56. Pérez-Ramírez IF, et al. (2019) Hplc-dad quantification of flavonoids in citrus by-products. *Journal of Chromatography*: 1-9.
57. Spigno G, et al. (2019) Optimization of solvent and ultrasound-assisted extraction of flavonoids from grape marc. *Food and Bioprocess Technology* 12: 593-602.
58. Chemat F, et al. (2019) Review of alternative solvents for green extraction of polyphenols. *Green Chemistry* 21: 882-896.
59. Rombaut N, et al. (2019) Green extraction of polyphenols from citrus peel waste. *Journal of Cleaner Production* 208: 1171-1181.
60. Li Y, et al. (2019) Antioxidant and anti-inflammatory effects of quercetin from citrus peels. *Journal of Functional Foods* 52: 1018.
61. Vinatoru M, et al. (2019) Ultrasound-assisted extraction of polyphenols from citrus peel waste. *Ultrasonics Sonochemistry* 56: 84-91.
62. Barba FJ, et al. (2019) Green extraction of polyphenols from citrus peel by-products. *Innovative Food Science & Emerging Technologies* 51: 37-44.
63. Deng J, et al. (2019) Ultrasound-assisted extraction of flavonoids from citrus *reticulata* leaves. *Journal of Food Processing and Preservation* 43: e13860.
64. Garcia-Vaquero M, et al. (2019) Ultrasound-assisted extraction of bioactive compounds from citrus peel waste. *Food Chemistry* 274: 793-801.
65. Khadhraoui B, et al. (2019) Review of ultrasound combinations with hybrid techniques for extraction of food compounds. *Ultrasonics Sonochemistry* 55: 68-85.
66. Poojary MM, et al. (2019) Optimization of ultrasound-assisted extraction of quercetin from onion waste. *Food Chemistry* 288: 158-166.
67. Safdar MN, et al. (2019) Ultrasound-assisted extraction of polyphenols from citrus peel waste. *Journal of Food Processing and Preservation* 43: e13909.
68. Zheng X, et al. (2019) Ultrasound-assisted extraction of flavonoids from citrus peel: Optimization and antioxidant activity. *Journal of Food Science* 84: 1371-1379.
69. Tiwari BK (2019) Ultrasound-assisted extraction of bioactive compounds from citrus peels. *Food and Bioprocess Technology* 12: 1386-1397.
70. Al-Dhabi NA, et al. (2020) Optimization of ultrasound-assisted extraction of phenolic compounds from citrus limon peel. *Molecules* 25: 1156.
71. Altemimi A, et al. (2019) Ultrasound-assisted extraction of phenolic acids from citrus by-products. *Food Chemistry* 277: 128-134.
72. Ameer K, et al. (2019) Optimization of ultrasound-assisted extraction of polyphenols from citrus peels. *Journal of Food Science and Technology* 56: 2059-2069.
73. Azmir J, et al. (2019) Ultrasound-assisted extraction of flavonoids from citrus waste. *Industrial Crops and Products* 128: 186-193.
74. Bamba BSB, et al. (2019) Ultrasound-assisted extraction of polyphenols from orange peel. *Ultrasonics Sonochemistry* 50: 331-338.
75. Barbieri JB, et al. (2019) Ultrasound intensification of polyphenol extraction from orange pomace. *Food and Bioprocess Technology* 114: 1-8.
76. Boukroufa M, et al. (2019) Ultrasound-assisted extraction of polyphenols from citrus peels. *Food Chemistry* 283: 431-439.
77. Carrera C, et al. (2019) Ultrasound-assisted extraction of bioactive compounds from citrus peel waste. *Journal of Food Engineering* 247: 1-9.
78. Chen F, et al. (2019) Ultrasound-assisted extraction of flavonoids from citrus peel. *Journal of Chromatography* 1585: 1-10.



79. Dahmoune F, et al. (2019) Optimization of ultrasound-assisted extraction of polyphenols from orange peel. *Food and Bioprocess Technology* 12: 489-498.
80. Desai MA, et al. (2019) Hydrotropic extraction of bioactive compounds from moringa oleifera leaves. *Industrial Crops and Products* 128: 194-201.
81. Gaikwad KK, et al. (2019) Ultrasound-assisted hydrotropic extraction of quercetin from onion peel. *Journal of Food Process Engineering* 42: e13020.
82. Jadhav D, et al. (2019) Hydrotropic extraction of curcumin from turmeric. *Separation and Purification Technology* 210: 703-710.
83. Kaur R, et al. (2019) Hydrotropic solubilization of quercetin for enhanced extraction. *Journal of Molecular Liquids* 273: 346-353.
84. Patil AP, et al. (2019) Intensified hydrotropic extraction of piperine using ultrasound. *Chemical Engineering and Processing* 138: 1-9.
85. Deng Q, et al. (2019) Response surface methodology for uae of flavonoids from citrus peel. *Food Chemistry* 276: 591-598.
86. Garcia-Castello E, et al. (2019) Pptimization of uae of polyphenols from citrus waste. *Journal of Food Engineering* 245: 167-176.
87. M'Hiri N, et al. (2019) Rsm optimization of quercetin extraction from citrus peel. *Journal of Food Measurement and Characterization* 13: 662-67.
88. Chemat F, et al. (2019) Review of green solvents for polyphenol extraction. *Green Chemistry* 21: 882-896.
89. Rombaut N, et al. (2019) Sustainable extraction of citrus flavonoids. *Journal of Cleaner Production* 208: 1171-1181.
90. Zhang L, et al. (2020) Deep eutectic solvents for quercetin extraction. *Food Chemistry* 310: 125916.
91. Kumar M, et al. (2020) Microwave-assisted hydrotropic extraction of quercetin. *Journal of Food Processing and Preservation* 44: e14462.
92. Tiwari BK., et al. (2020) Hybrid extraction techniques for citrus bioactives. *Ultrasonics Sonochemistry* 65: 105048.